

A “sodium trap” based on benzo-15-crown-5 with an exocyclic *N*-(thiophosphoryl)thiourea moiety

Felix D. Sokolov ^a, Damir A. Safin ^{a,*}, Maria G. Babashkina ^a, Nail G. Zabiroy ^a,
 Vasiliy V. Brusko ^a, Nikolay A. Mironov ^a, Dmitry B. Krivolapov ^b, Igor A. Litvinov ^b,
 Rafael A. Cherkasov ^a, Boris N. Solomonov ^a

^a Department of Chemistry, Kazan State University, Kremlevskaya Street 18, 420008 Kazan, Russia

^b Arbuzov Institute of Organic and Physical Chemistry, Arbuzov Street 8, 420088 Kazan, Russia

Received 12 October 2006; accepted 7 December 2006

Available online 22 December 2006

Abstract

For *N*-(thio)phosphorylthioureas of the common formula RC(S)NHP(X)(OiPr)₂ **HL**^I (R = *N*-(4'-aminobenzo-15-crown-5), X = S), **HL**^{II} (R = *N*-(4'-aminobenzo-15-crown-5), X = O), **HL**^{III} (R = PhNH, X = S), **HL**^{IV} (R = PhNH, X = O), and (*N,N'*-bis-[C(S)NHP(S)(OiPr)₂]₂-1,10-diaza-18-crown-6) **H₂L**^V, salts **LiL**^{I,III,IV}, **NaL**^{I-IV}, **KL**^{I-IV}, **M₂L**^V (M = Li⁺, Na⁺, K⁺), **Ba(L**^{I,III,IV})₂, and **BaL**^V have been synthesized and investigated. Compounds **NaL**^{I,III} quantitatively drop out as a deposit in ethanol medium, allowing the separation of Na⁺ and K⁺ cations. This effect is not displayed for the other compounds. The crystal structures of **HL**^{III} and the solvate of the composition [**K(Me₂CO)L**^{III}] have been investigated by X-ray crystallography.

© 2006 Elsevier Ltd. All rights reserved.

Keywords: Chelates; Coordination modes; Crown-ethers; Crystal structures; *N*-thioacylamido(thio)phosphates

1. Introduction

Crown-ethers and azamacrocycles modified by exocyclic groups are widely applied as complexing agents, selective to cations of alkaline, earth-alkaline, and d-metals [1,2], lanthanides and actinides [3], and Tl(I) [4]. However, in the literature there is no data about direct (not-extraction) highly effective separation of alkaline metal cations using crown compounds.

On the formation of complexes of alkaline metal salts [M⁺A[−]] with crown-ethers, as a rule, there is an increase in solubility of the formed adduct [M(crown)]⁺A[−] in an organic phase [5]. However, a combination of crown-ether and negative charged chelating units in a ligand molecule can lead to interesting cooperative effects.

Herein we present data about the structures and properties of *N*-(thio)phosphorylthioureas of the common formula RC(S)NHP(X)(OiPr)₂ (**HL**^{I-IV}) and *N,N'*-bis-[C(S)NHP(S)(OiPr)₂]₂-1,10-diaza-18-crown-6 (**H₂L**^V) (Scheme 1), and their salts with Li⁺, Na⁺, K⁺, Ba²⁺.

It is known [6] that *N*-(thio)phosphorylthioureas in aqueous alcohol medium are weak acids: **HL**^{III} p*K* 8.1 (96% aq. EtOH) [7], 8.28 (96% aq. *i*PrOH) [8]; **HL**^{IV} p*K* 6.93 (96% aq. *i*PrOH) [8]; **H₂L**^V p*K*₁ = 9.2, p*K*₂ = 10.2 (96% aq. EtOH) [7]. These compounds are able to form salts with cations of alkaline metals. There is no information on the crystal structures of salts of *N*-(thio)phosphorylthioureas in the literature. However, the structures of some close analogues of these compounds – salts of *N*-(thio)phosphorylamides and thioamides, containing the fragment C(X)NHP(Y)R₂ (X, Y = O, S), have already been investigated. Compounds of the formula [M(CCl₃C(O)NP(O)(OMe)₂)] as crystals form dimers (M = Na⁺) or polymeric chains M = Rb⁺ [9]. Potassium salts of thioamides

* Corresponding author. Tel./fax: +7 843 2543734.

E-mail addresses: felix.sokolov@ksu.ru (F.D. Sokolov), damir.safin@ksu.ru (D.A. Safin).